

Solution of the quasipositivity problem in the braid group

A.Bentalha

1st February 2008

Introduction Let B_{n+1} be the group of braids with $n + 1$ strings. It has a presentation ,

$$(\Theta) : B_{n+1} = [\sigma_1, \dots, \sigma_n \mid \sigma_j \sigma_k = \sigma_k \sigma_j \text{ for } |j - k| > 1, \sigma_j \sigma_k \sigma_j = \sigma_k \sigma_j \sigma_k \text{ for } |j - k| = 1]$$

A braid b in B_{n+1} is called *quasipositive* if there are braids $w_1, \dots, w_k \in B_{n+1}$ such that

$$(\Upsilon) : b = (w_1 \sigma_{i_1} w_1^{-1})(w_2 \sigma_{i_2} w_2^{-1}) \dots (w_k \sigma_{i_k} w_k^{-1})$$

each w_j can be supposed positive.

The purpose of this paper is to solve *the quasipositivity problem* in B_{n+1} , i.e. to find an algorithm which decides if a given braid is quasipositive or not. This result can be used in the 16th Hilbert problem, which deals with the classification of real algebraic curves.

We refer the reader to *Orevkov* [7],[8], *Rudolph* [9],[10] for more motivations. Our algorithm is based on the ideas of Garside, see *Garside* [5].

It is based on Theorem 1 below which was conjectured by S. Orevkov.

Let us denote by B_{n+1}^+ the *monoid of positive braids*, this is the monoid, (semigroup with the unit) defined by the presentation (Θ) , see *Birman* [1], *Garside* [5] for details.

It is known by *Garside* that B_{n+1}^+ coincides with the submonoid of B_{n+1} generated by $\sigma_1, \dots, \sigma_n$.

This means that two positive words in $\sigma_1, \dots, \sigma_n$ represent the same element of B_{n+1} if and only if one can pass from one to another using only *positive* relations.

Moreover these *positive* relations don't change the number of generators so we can define on B_{n+1}^+ a unique length function giving this number.

We say that a (left) divides b (a right divides b) and we note $a \prec b$ ($b \succ a$) if there exists c satisfying $b = ac$ ($b = ca$).

Let Δ be *the Garside element* of B_{n+1} :

$$\Delta = (\sigma_1 \sigma_2 \dots \sigma_n)(\sigma_1 \sigma_2 \dots \sigma_{n-1}) \dots (\sigma_1 \sigma_2) \sigma_1$$

This element is "almost central", i.e. $\Delta\sigma_k = \sigma_{n+1-k}\Delta$ for $1 \leq k \leq n$. Let us denote σ_{n+1-k} by $\bar{\sigma}_k$. We then obtain, for any word a in B_{n+1} : if s even $\Delta^s a = a\Delta^s$ and if s odd, $\Delta^s a = \bar{a}\Delta^s$; in the next, we note $\Delta^s a = \bar{a}\Delta^s$ when we don't know if s is even or not.

This implies that a positive braid b is right divisible by Δ in B_{n+1}^+ if and only if it is (left) divisible and we shall say in this case that b is *divisible by Δ* . We call by simple element, every positive word which divides Δ .

Any braid can be represented in the form $\Delta^{-r}w$ with an integer r and a positive braid w which is not divisible by Δ .

Moreover, there is an effective algorithm which allows to compute such a representation, see *Garside [5], Morton, El Rifai [6]* and we give here a Garside's lemma useful for the next.

Lemma[Garside] 0 1 *We give positive words X, Y such as $\sigma_i X = \sigma_j Y$ for generators σ_i, σ_j .*

If $i = j$, then $X = Y$.

If $|i - j| = 1$, then there exists Z such as $X = \sigma_j \sigma_i Z$ and $Y = \sigma_i \sigma_j Z$.

If $2 \leq |i - j|$, then there exists Z such as $X = \sigma_j Z$ and $Y = \sigma_i Z$.

In the next paragraph, we give series of results and in the following one their proofs. In the last one we give a computational algorithm and we study its complexity.

Statement of the result

Definition 1 1 *Let $w \in B_{n+1}^+$, $w = u\sigma v$. We note*

$$w = u\sigma v \xrightarrow{1} uv$$

to say that we pass from w to uv by eliminating one generator. This last is said to be a 1-sequence. So for any X in B_{n+1}^+ , the k -sequence

$$X \xrightarrow{k} Y$$

means that we pass from X to Y by eliminating k generators, i.e there exists $k + 1$ positive words L_j with $0 \leq j \leq k$ such as

$$L_k = X$$

$$L_{j+1} \xrightarrow{1} L_j$$

$$L_0 = Y$$

REMARK:

$X \xrightarrow{k} Y$ represents one of the way to reduce X to Y . When we have to be more precise, we write

$$X = X_k \delta_{i_k} Y_k \xrightarrow{1} X_k Y_k = X_{k-1} \delta_{i_{k-1}} Y_{k-1} \xrightarrow{k-2} X_1 \delta_{i_1} Y_1 \xrightarrow{1} X_1 Y_1 = Y.$$

Moreover, if there exist others $(X'_j, Y'_j, \delta'_{i_j})$ reducing X to Y , we say that the two k -sequences are equivalent.

EXAMPLE:

We note 1 for σ_1 , 1^{-1} for σ_1^{-1} , 2 for σ_2 and so on...

Let the word 2131 be in B_4 then

$$2132 \xrightarrow{1} 212 = 121 \xrightarrow{1} 11$$

that is

$$2132 \xrightarrow{2} 11$$

We see in this example that we can't delete two generators straight away of the initial word; we have to transform 212 into 121.

The k -sequence $X \xrightarrow{k} Y$ is more general than deleting k generators at the same time and in the same word X .

We can summarise the results of the whole paper, by:

Fundamental Result 1 *For positive words X, Y , with $\text{length}(X) = \text{length}(Y) + k$, the following are equivalent:*

1. $Y^{-1}X$ is quasipositive
2. $X \xrightarrow{k} Y$
3. $X d_1 d_2 d_3 \dots d_k = Y d_1 \sigma_{i_1} d_2 \sigma_{i_2} d_3 \dots \sigma_{i_{k-1}} d_k \sigma_{i_k}$ for positive d_j and generators σ_{i_j}

The d_j 's are defined in the next. In this case, one quasipositive form of $Y^{-1}X$ is

$$Y^{-1}X = (d_1 d_2 \dots d_k)^{-1} d_1 \sigma_{i_1} d_2 \sigma_{i_2} d_3 \dots \sigma_{i_{k-1}} d_k \sigma_{i_k}$$

Moreover, we have a finite number of quasipositive forms for $Y^{-1}X$; each form corresponds to a way to reduce X to Y , by $X \xrightarrow{k} Y$, and theses last are finite.

Lemma 1 1 *For positive words A, X, Y , and a positive integer k , the k -sequence $AX \xrightarrow{k} AY$ is equivalent to the other one $X \xrightarrow{k} Y$*

Theorem 1 1 *We give $b = \Delta^{-r} b_1 \dots b_s = N^{-1} P$ written in its normal form, with algebraic length k , then b is quasipositive if and only if $P \xrightarrow{k} N$.*

PROOF:

Let b be quasipositive, $b = \Delta^{-r} b_+ :$ by definition $b = w_1 \sigma_{i_1} w_1^{-1} \dots w_k \sigma_{i_k} w_k^{-1}$ for positive w_i . Each w_i written in its normal form gives $w_i = \Delta^{m_i} w_{i+}$ and also $w_i^{-1} = \Delta^{-p_i} v_{i+}$ for m_i, p_i positive integer. Moreover $w_i w_i^{-1} = 1$ is equivalent to $\Delta^{m_i} w_{i+} \Delta^{-p_i} v_{i+} = 1$, $\Delta^{m_i - p_i} w_{i+} v_{i+} = 1$; finally $w_{i+} v_{i+} = \Delta^{p_i - m_i}$.

We can write b as $b = \Delta^{m_1-p_1+\dots+m_k-p_k} w_{1+} \tilde{\sigma}_{i_1} v_{1+} \dots w_{k+} \tilde{\sigma}_{i_k} v_{k+}$, and we note w for $w_{1+} \tilde{\sigma}_{i_1} v_{1+} \dots w_{k+} \tilde{\sigma}_{i_k} v_{k+}$ and $m-p$ for $m_1-p_1+\dots+m_k-p_k$.

If we cancel in w all the generators σ_{i_j} , w becomes $w_{1+} v_{1+} \dots w_{k+} v_{k+}$ which is equal to $\Delta^{p_1-m_1+\dots+p_k-m_k} = \Delta^{p-m}$. In other words,

$$w \xrightarrow{k} \Delta^{p_1-m_1+\dots+p_k-m_k}$$

Futhermore, b in its normal form satisfies $w = \Delta^u b_+$, for an integer u .

As $b = \Delta^{-r} b_+$ and $b = \Delta^{m-p} w$, we obtain $\Delta^{m-p} w = \Delta^{-r} b_+$, $w = \Delta^{p-m-r} b_+$.

Finally $r+u=p-m$ and the sequence becomes

$$w = \Delta^u b_+ \xrightarrow{k} \Delta^{p-m} = \Delta^u \Delta^r$$

By lemmas above, $AX \xrightarrow{k} AY$ beeing equivalent to $X \xrightarrow{k} Y$, we obtain here, by simplifying by Δ^u ,

$$b_+ \xrightarrow{k} \Delta^r$$

Moreover $b = N^{-1}P$ satisfies by definition $RP = b_+$ and $RN = \Delta^r$ for a positive R : the sequence above becomes

$$b_+ = RP \xrightarrow{k} \Delta^r = RN \text{ which is equivalent to}$$

$$P \xrightarrow{k} N$$

Conversely, if $b = N^{-1}P$ satisfies $P \xrightarrow{k} N$, we can write $P = \sigma_{i_1} \dots \sigma_{i_k} N$ and obviously $b = N^{-1} \sigma_{i_1} \dots \sigma_{i_k} N$ is quasipositive.

EXAMPLE:

In B_4 , $b = 212^{-1}2^{-1}32$ is quasipositive, with $k = 2$: $b = 1^{-1}1^{-1}2132$ in its normal form; we have

$$P = 2132 \xrightarrow{2} N = 11; \text{ indeed}$$

$$2132 \xrightarrow{1} 212 = 121 \xrightarrow{1} 11$$

Theorem 11 gives us a therotical way to decide whether a word is quasipositive or not. We now give some results, proven in the following, which lead us to a computational algorithm.

Dealing with a k -sequence $X \xrightarrow{k} Y$, we saw that the k generators deleted are not necessary deleted at the same time; but we see in the next that we can trasform any sequence into one of the form

$$A_1 \sigma_{i_1} A_2 \dots A_k \sigma_{i_k} A_{k+1} \xrightarrow{k} A_1 A_2 \dots A_k A_{k+1}$$

where the k generators can be deleted in the same word and at the same time.

Lemma 2 1 $b = \Delta^{-r}b_+$ with algebraic length k is quasipositive if and only if there exist $k + 1$ positive words $A_1, \dots, A_{k+1}$ and k generators $\sigma_{i_1}, \dots, \sigma_{i_k}$ such as

$$b_+ = A_1\sigma_{i_1}A_2\sigma_{i_2}\dots A_k\sigma_{i_k}A_{k+1}$$

$$\Delta^{-r} = A_1A_2\dots A_kA_{k+1}$$

This lemma gives a stronger result than $b_+ \xrightarrow{k} \Delta^{-r}$. Unfortunately, we can't obtain a similar version for $P \xrightarrow{k} N$. But we have a slightly different one, which will give the main tool for an algorithm.

Lemma 3 1 We give $b = \Delta^{-r}b_+ = \Delta^{-r}b_1b_2\dots b_r\dots b_s$ with algebraic length k . We define for $0 \leq j \leq r$,

$$P_j = b_{j+1}\dots b_{s-r+j}$$

and N_j such as

$$b_1\dots b_jN_jb_{s-r+j+1}\dots b_s = \Delta^r$$

b is quasipositive if and only if for one j , $0 \leq j \leq r$, there exist k positive words $A_2, \dots, A_{k+1}$ (here $A_1 = 1$) and k generators $\sigma_{i_1}, \dots, \sigma_{i_k}$ such as

$$P_j = \sigma_{i_1}A_2\sigma_{i_2}\dots A_k\sigma_{i_k}A_{k+1}$$

$$N_j = A_2\dots A_kA_{k+1}$$

In this case, we output the quasipositive form of b : we set $T_j = (b_{s-r+j+1}\dots b_s)$ and

$$T_j b T_j^{-1} = N_j^{-1} P_j \text{ so}$$

$$b = T_j^{-1}[(A_2\dots A_kA_{k+1})^{-1}(\sigma_{i_1}A_2\sigma_{i_2}\dots A_k\sigma_{i_k}A_{k+1})]T_j$$

Technical proofs The next lemma is a classical known result.

Lemma 4 1 lgcd (left greatest common divisor) is distributive.

Let R, A, B be words in B_{n+1} , $\text{lgcd}(RA, RB) = R \text{lgcd}(A, B)$.

(resp. rgcd (right greatest common divisor) is distributive, $\text{rgcd}(AR, BR) = \text{rgcd}(A, B)R$).

PROOF:

$A = \text{lgcd} (A, B)A'$ and $B = \text{lgcd} (A, B)B'$ by definition of lgcd , for positive A', B' .

So $RA = R \text{lcd} (A, B)A'$ and $RB = R \text{lgcd} (A, B)B'$, that is

$R \text{lgcd} (A, B) \prec \text{lgcd} (RA, RB)$, $\text{lgcd} (RA, RB)$ beeing the left greatest common divisor.

Let U be such as $\text{lgcd} (RA, RB) = R \text{lgcd} (A, B)U$.

Then $\text{lgcd} (RA, RB) = R \text{lgcd} (A, B)U \prec RA = R \text{lgcd} (A, B)A'$ gives

$R \text{lgcd} (A, B)U \prec R \text{lgcd} (A, B)A'$, so $U \prec A'$ and in the same way $U \prec B'$.

Finally, $\text{lgcd} (A, B)U$ divides $\text{lgcd} (A, B)A' = A$ and also B ;

but $\text{lgcd} (A, B)$ beeing the left greatest common divisor of A and B , $U = 1$ and we have the result,

$\text{lgcd} (RA, RB) = R \text{lgcd} (A, B)$.

The next one, very important for the next, already appeared in some work of Juan Gonzalez-Meneses.

Lemma 5 1 *Let w be a word in B_{n+1}^+ and σ a generator.*

We note $D_w^l(\sigma) = (X \text{ positive such as } wX \prec \sigma wX)$.

In this case $wX\sigma' = \sigma wX$ for a generator σ' .

$D_w^l(\sigma)$ is not empty and has a minimal element for left divisibility, called $d_w^l(\sigma)$, this last divides all $X \in D_w^l(\sigma)$.

(We have same results for right side: $d_w^r(\sigma)$ minimal for right divisibility in $D_w^r(\sigma) = (Y \text{ positive such that } Yw\sigma \succ Yw)$).

PROOF:

Let w be written in its left normal form,

$$w = \Delta^{-r}b_1 \dots b_s$$

with $-r$ integer and let a_i be such as $b_i a_i = \Delta$ for $1 \leq i \leq s$.

We have $b_1 \dots b_{s-1} b_s a_s a_{s-1}^{-1} \dots \tilde{a}_1 = \Delta^s$ and if we note $X = a_s a_{s-1}^{-1} \dots \tilde{a}_1$ then X satisfies $\sigma wX = \sigma \Delta^{-r} b_1 \dots b_s a_s a_{s-1}^{-1} \dots \tilde{a}_1 = \sigma \Delta^{-r+s} = \Delta^{-r+s} \tilde{\sigma}$ that is

$$\sigma wX = wX\tilde{\sigma}$$

and $X \in D_w^l(\sigma)$.

Let's now prove that $D_w^l(\sigma)$ has a minimal element: if X and Z are in $D_w^l(\sigma)$, also is $\text{lgcd}(X, Z)$, indeed

$$\sigma wX = wX\sigma'$$

$$\sigma wZ = wZ\sigma''$$

for generators σ', σ'' .

Then $\text{lgcd}(\sigma w X, \sigma w Z) = \text{lgcd}(w X \sigma', w Z \sigma'')$ and by Lemma 31 on the distributivity of lgcd , $\sigma w \text{lgcd}(X, Z) = w \text{lgcd}(X \sigma', Z \sigma'')$.

But $\text{lgcd}(X, Z) \prec X \prec X \sigma'$ and $\text{lgcd}(X, Z) \prec Z \prec Z \sigma''$ so

$\text{lgcd}(X, Z) \prec \text{lgcd}(X \sigma', Z \sigma'')$ and $w \text{lgcd}(X, Z)u = w \text{lgcd}(X \sigma', Z \sigma'')$ for a positive u .

If we replace this in what before, $\sigma w \text{lgcd}(X, Z) = w \text{lgcd}(X, Z)u$.

Now, dealing with positive words, the algebraic length of the two factors in the equality must be equal, so u is a generator, $u = \beta$

Finally $\sigma w \text{lgcd}(X, Z) = w \text{lgcd}(X, Z)u$ gives $\sigma w \text{lgcd}(X, Z) = w \text{lgcd}(X, Z)\beta$ so $\text{lgcd}(X, Z)$ satisfies the same properties as X, Z ,

$\text{lgcd}(X, Y) \in D_w^l(\sigma)$.

And $d_w^l(\sigma)$ can be taken as being the common lgcd of all words in $D_w^l(\sigma)$.

Lemma 6 1 *Let w be a positive word, σ a generator and $d_w^l(\sigma)$ as above:*

$\sigma w d_w^l(\sigma) = w d_w^l(\sigma) \sigma'$.

If f and h are positive words such as

$$\sigma w f = w h$$

then, $d_w^l(\sigma) \prec f$, $d_w^l(\sigma) \prec h$ and if $f = d_w^l(\sigma)x$, $h = d_w^l(\sigma)\sigma'x$ for a positive x .

(We have same results for right divisibility).

PROOF:

We use Garside's Lemma to prove it when w is a generator: $w = \beta$

if $\sigma = \beta$ then $\sigma\beta = \beta\sigma$ and $d_w^l(\sigma) = 1$, $d_w^l(\sigma)$ obviously divides f as above

if σ and β commutes then again $\sigma\beta = \beta\sigma$ and $d_w^l(\sigma) = 1$ as above

if $\sigma\beta\sigma = \beta\sigma\beta$, then $d_w^l(\sigma) = \sigma$ and we know:

$\sigma\beta f = \beta h$ gives $f = \sigma x$ and $h = \sigma\beta x$ for a positive x which gives

$d_w^l(\sigma) \prec (f, h)$, $f = d_w^l(\sigma)x$, $h = d_w^l(\sigma)\sigma'x$, $\sigma' = \beta$.

(we have same results for right divisibility)

So let's write now the proof in the general case:

$$\sigma w d_w^l(\sigma) = w d_w^l(\sigma) \sigma'$$

$$\sigma w f = w h$$

We have $\text{lgcd}(\sigma w d_w^l(\sigma), \sigma w f) = \text{lgcd}(w d_w^l(\sigma) \sigma', w h)$ which is equivalent to $\sigma w \text{lgcd}(d_w^l(\sigma), f) = w \text{lgcd}(d_w^l(\sigma) \sigma', h)$.

If we note $a = \text{lgcd}(d_w^l(\sigma), f)$ and $b = \text{lgcd}(d_w^l(\sigma) \sigma', h)$:

$au = d_w^l(\sigma)$ for a positive u and $\sigma w a = w b$ gives $\sigma w a u = w b u$, that is $\sigma w d_w^l(\sigma) = w b u$ but $\sigma w d_w^l(\sigma) = w d_w^l(\sigma) \sigma'$ so

$$b u = d_w^l(\sigma) \sigma'$$

If u no equals to 1 then $u = v\delta$ for a generator δ :

$au = d_w^l(\sigma)$ gives $au\sigma' = bu$ that is $av\delta\sigma' = bv\delta$.

The word δ 'pseudocommutes' with σ' and $d_\delta^r(\sigma')$ satisfies $d_\delta^r(\sigma')\delta\sigma' = \sigma''d_\delta^r(\sigma')\delta$; there exists y positive, as a consequence of Lemma 5 1, with

$$av = yd_\delta^r(\sigma')$$

$$bv = y\sigma''d_\delta^r(\sigma')$$

Finally,

$$\sigma wd_w^l(\sigma) = wd_w^l(\sigma)\sigma'$$

$$\sigma wau = wbu$$

$$\sigma wav\delta = wbv\delta$$

$$\sigma wyd_\delta^r(\sigma')\delta = wy\sigma''d_\delta^r(\sigma')\delta$$

$$\sigma wy = wy\sigma''$$

and y satisfies the same propriety as $d_w^l(\sigma)$, y belongs to $D_w^l(\sigma)$.

But $yd_\delta^r(\sigma')\delta = d$ and y is strictly smaller than $d_w^l(\sigma)$; which is a contradiction of the minimality of d_w^l in $D_w^l(\sigma)$.

As all this results from u not equal to 1 ,we obtain $u = 1$ and

$$\text{lgcd}(d_w^l(\sigma), f) = a = au = d_w^l(\sigma)$$

$$\text{lgcd}(d_w^l(\sigma)\sigma', h) = b = bu = d_w^l(\sigma)\sigma'$$

$$\text{so } d_w^l(\sigma) \prec f, d_w^l(\sigma)\sigma' \prec h;$$

if we put $f = d_w^l(\sigma)x$, then $\sigma wf = wh$ gives

$$\sigma wd_w^l(\sigma)x = wh$$

$$wd_w^l(\sigma)\sigma'x = wh$$

and $h = d_w^l(\sigma)\sigma'x$, which is what we are looking for.

Lemma 7 1 *Let $X \xrightarrow{k} Y$ be a k -sequence, then there exists k positive words $d_1, \dots, d_k$ and k generators $\sigma_{i_1}, \dots, \sigma_{i_k}$ such as*

$$Xd_1d_2\dots d_k = Yd_1\sigma_{i_1}d_2\sigma_{i_2}\dots d_k\sigma_{i_k}$$

PROOF:

$X \xrightarrow{k} Y$ means there exist $X_j, Y_j, \delta_{i_j}, 1 \leq j \leq k$ satisfying

$$X = X_k\delta_{i_k}Y_k \xrightarrow{1} X_kY_k = X_{k-1}\delta_{i_{k-1}}Y_{k-1} \xrightarrow{k-2} X_2Y_2 = X_1\delta_{i_1}Y_1 \xrightarrow{1} Y = X_1Y_1$$

We now consider :

$$d_1 = d_{Y_1}{}^r(\delta_{i_1}) \text{ which satisfies by Lemma 5 1, } \delta_{i_1} Y_1 d_1 = Y_1 d_1 \delta'_{i_1}$$

$$d_2 = d_{Y_2 d_1}{}^r(\delta_{i_2}) \text{ such as } \delta_{i_2} Y_2 d_1 d_2 = Y_2 d_1 d_2 \delta'_{i_2}$$

$$d_j = d_{Y_j d_1 d_2 \dots d_{j-1}}{}^r(\delta_{i_j}) \text{ such as } \delta_{i_j} Y_j d_1 d_2 \dots d_{j-1} d_j = Y_j d_1 d_2 \dots d_{j-1} d_j \delta'_{i_j}$$

$$d_k = d_{Y_k d_1 d_2 \dots d_{k-1}}{}^r(\delta_{i_k}) \text{ such as } \delta_{i_k} Y_k d_1 d_2 \dots d_{k-1} d_k = Y_k d_1 d_2 \dots d_{k-1} d_k \delta'_{i_k}$$

By definition of theses d_j , we have by multiplying the sequence above by

$$d_1 d_2 \dots d_k:$$

for the first line , $Y d_1 d_2 \dots d_k$

for the second one, $X_2 Y_2 d_1 d_2 \dots d_k = X_1 \delta_{i_1} Y_1 d_1 d_2 \dots d_k = X_1 Y_1 d_1 \delta'_{i_1} d_2 \dots d_k =$

$Y d_1 \delta'_{i_1} d_2 \dots d_k$; moreover this equality gives $X_2 Y_2 d_1 = Y d_1 \delta'_{i_1}$

for the third one, $X_3 Y_3 d_1 d_2 \dots d_k = X_2 \delta_{i_2} Y_2 d_1 d_2 \dots d_k = X_2 Y_2 d_1 d_2 \delta'_{i_2} d_3 \dots d_k =$

$Y d_1 \delta'_{i_1} d_2 \delta'_{i_2} d_3 \dots d_k$ by what above; and we also obtain $X_3 Y_3 d_1 d_2 = Y d_1 \delta'_{i_1} d_2 \delta'_{i_2}$

for the k th one, $X_k Y_k d_1 d_2 \dots d_k = X_{k-1} \delta_{i_{k-1}} Y_{k-1} d_1 d_2 \dots d_k = X_{k-1} Y_{k-1} d_1 d_2 \dots d_{k-1} \delta'_{i_{k-1}} d_k =$

$Y d_1 \delta'_{i_1} d_2 \delta'_{i_2} d_3 \delta'_{i_3} \dots d_{k-1} \delta'_{i_{k-1}} d_k$ by induction,

and in the same way, the final result $X d_1 d_2 \dots d_k = Y d_1 \delta'_{i_1} d_2 \delta'_{i_2} d_3 \delta'_{i_3} \dots d_{k-1} \delta'_{i_{k-1}} d_k \delta'_{i_k}$

In the next Lemma, we see that within the condition

$$X d_1 d_2 \dots d_k = Y d_1 \sigma_{i_1} d_2 \sigma_{i_2} \dots d_k \sigma_{i_k}, \text{ our } d_j \text{ are minimal.}$$

Lemma 8 1 *Let $X = X_k \delta_{i_k} Y_k \xrightarrow{1} X_k Y_k = X_{k-1} \delta_{i_{k-1}} Y_{k-1} \xrightarrow{k-1} X_1 Y_1 = Y$ be a k -sequence.*

If there exists k positive words $A_1, \dots, A_k$ and k generators $\sigma_{i_1}, \dots, \sigma_{i_k}$ such as $X A_1 A_2 \dots A_k = Y A_1 \sigma_{i_1} A_2 \sigma_{i_2} \dots A_k \sigma_{i_k}$, then we can find an equivalent k -sequence

$$X = X'_k \delta'_{i_k} Y'_k = L_k \xrightarrow{1} X'_k Y'_k = X'_{k-1} \delta'_{i_{k-1}} Y'_{k-1} = L_{k-1} \xrightarrow{k-1} X'_1 Y'_1 = Y = L_0 \text{ such as}$$

$$L_j A_1 \dots A_j = Y A_1 \sigma_{i_1} A_2 \sigma_{i_2} \dots A_j \sigma_{i_j}$$

PROOF:

By induction on $l = \text{length}(Y)$.

$l = 0$. Which is equivalent to $Y = 1$. Let $X \xrightarrow{k} 1$ be a k -sequence with $X A_1 A_2 \dots A_k = A_1 \sigma_{i_1} A_2 \sigma_{i_2} \dots A_k \sigma_{i_k}$. We use now an induction on k .

$k = 1$ gives $X = X_1 \delta_{i_1} Y_1 \xrightarrow{1} 1$ and $X A_1 = A_1 \sigma_{i_1}$. Obviously $L_1 = X$ satisfies $L_1 A_1 = A_1 \sigma_{i_1}$, so the result.

We suppose the result true for any $(k-1)$ -sequence. With the previous datas, we set

$$U = A_1 \sigma_{i_1} A_2 \sigma_{i_2} \dots A_{k-1} \sigma_{i_{k-1}} \cdot (A_1 \dots A_{k-1})^{(-1)}$$

Which gives $U A_1 \dots A_{k-1} = Y A_1 \sigma_{i_1} A_2 \sigma_{i_2} \dots A_{k-1} \sigma_{i_{k-1}}$ and by replacing in $X A_1 A_2 \dots A_k = A_1 \sigma_{i_1} A_2 \sigma_{i_2} \dots A_k \sigma_{i_k}$, we obtain $X A_1 A_2 \dots A_k = U A_1 A_2 \dots A_k \sigma_{i_k}$.

U is positive and by using D minimal, as in Lemma 5 1, such as $DA_1A_2...A_k\sigma_{i_k} = \sigma'_{i_k}DA_1A_2...A_k$,

$$\begin{aligned} X &= E\sigma'_{i_k}D \\ U &= ED \end{aligned}$$

for E positive.

Moreover, $\text{length}(U) = k-1$ gives $U \xrightarrow{k-1} 1$. As, $UA_1...A_{k-1} = YA_1\sigma_{i_1}A_2\sigma_{i_2}...A_{k-1}\sigma_{i_{k-1}}$, the hypothesis of induction gives

$$U = L_{k-1} \xrightarrow{k-j} L_j \xrightarrow{j} L_0 = 1$$

with $L_jA_1...A_j = A_1\sigma_{i_1}A_2\sigma_{i_2}...A_j\sigma_{i_j}$. If we add $X = E\sigma'_{i_k}D \xrightarrow{1} U = ED$, we obtain a k -sequence with each line L_j satisfying the final result.

For any l . We suppose the result true for $l-1$. Let $X \xrightarrow{k} Y$ with $\text{length}(Y) = l$.

With $Y = Y'\delta$, we obtain a $(k+1)$ -sequence

$$X \xrightarrow{k} Y = Y'\delta \xrightarrow{1} Y'$$

By setting $1 = A_0$ and $\delta = \sigma_{i_0}$, the equality $XA_1A_2...A_k = YA_1\sigma_{i_1}A_2\sigma_{i_2}...A_k\sigma_{i_k}$ becomes $XA_0A_1A_2...A_k = Y'A_0\sigma_{i_0}A_1\sigma_{i_1}A_2\sigma_{i_2}...A_k\sigma_{i_k}$.

As $\text{length}(Y') = l-1$, we have by hypothesis of induction, an equivalent $(k+1)$ -sequence $X = L_k \xrightarrow{k-j} L_j \xrightarrow{j} L_0 \xrightarrow{1} L'_0 = Y'$. Each L_j satisfies $L_jA_0A_1...A_j = Y'A_0\sigma_{i_0}A_1\sigma_{i_1}A_2\sigma_{i_2}...A_j\sigma_{i_j}$. And L_0 is such as $L_0A_0 = Y'A_0\sigma_{i_0}$ which is $L_0 = Y$.

And the final result $X = L_k \xrightarrow{k-j} L_j \xrightarrow{j} L_0 = Y$, with $L_jA_0A_1...A_j = Y'A_0\sigma_{i_0}A_1\sigma_{i_1}A_2\sigma_{i_2}...A_j\sigma_{i_j}$ equivalent to $L_jA_1...A_j = YA_1\sigma_{i_1}A_2\sigma_{i_2}...A_j\sigma_{i_j}$.

Lemma 9 1 *Conversely, if there exists k positive words $A_1, ..., A_k$ and k generators $\sigma_{i_1}, ..., \sigma_{i_k}$ such as $XA_1A_2...A_k = YA_1\sigma_{i_1}A_2\sigma_{i_2}...A_k\sigma_{i_k}$, then*

$$X \xrightarrow{k} Y$$

PROOF:

By induction on $l = \text{length}(Y)$.

$l = 0$: which is equivalent to $Y = 1$.

$XA_1A_2...A_k = YA_1\sigma_{i_1}A_2\sigma_{i_2}...A_k\sigma_{i_k}$ becomes $XA_1A_2...A_k = A_1\sigma_{i_1}A_2\sigma_{i_2}...A_k\sigma_{i_k}$. The equality of length in the positive monoid gives $l(X) = k$ and obviously

$$X \xrightarrow{k} 1$$

Any l . For Y' with length $l - 1$ we suppose the result true: for any k integer and $XA_1A_2...A_k = YA_1\sigma_{i_1}A_2\sigma_{i_2}...A_k\sigma_{i_k}$ then $X \xrightarrow{k} Y$. We give now $Y = Y'\delta$ of length l satisfying $XA_1A_2...A_k = Y'\delta A_1\sigma_{i_1}A_2\sigma_{i_2}...A_k\sigma_{i_k}$ for positive A_i 's.

We consider $A_0 = 1$ and $\sigma_{i_0} = \delta$ to have a new equality $XA_0A_1A_2...A_k = Y'A_0\sigma_{i_0}A_1\sigma_{i_1}A_2\sigma_{i_2}...A_k\sigma_{i_k}$.

But $l(Y') = l - 1$, we can apply what before to obtain $X \xrightarrow{k+1} Y'$. Lemma 8 1 tells us that we have an equivalent k-sequence such as

$$X = L_k \xrightarrow{k-j} L_j \xrightarrow{j} L_0 \xrightarrow{1} L'_0 = Y' \text{ and } L_j A_0 A_1 ... A_j = Y' A_0 \sigma_{i_0} A_1 \sigma_{i_1} A_2 \sigma_{i_2} ... A_j \sigma_{i_j}.$$

In a particular case, $L_0 A_0 = Y' A_0 \sigma_{i_0}$ which is $L_0 = Y$.

And the final k-sequence $X \xrightarrow{k} L_0 = Y$.

We now give the proof of Lemma 1 1.

PROOF Lemma 1 1:

If $X \xrightarrow{k} Y$, obviously for any positive A , $AX \xrightarrow{k} AY$.

Conversely if we have $AX \xrightarrow{k} AY$, Lemma 7 1 gives $d_1, d_2, ..., d_k$ and $\sigma_{i_1}, \sigma_{i_2}, ..., \sigma_{i_k}$ such as

$$AXd_1d_2...d_k = AYd_1\sigma_{i_1}d_2\sigma_{i_2}...d_k\sigma_{i_k}$$

This last equality is equivalent to $Xd_1d_2...d_k = Yd_1\sigma_{i_1}d_2\sigma_{i_2}...d_k\sigma_{i_k}$ when simplifying by A .

And Lemma 9 1 finally gives $X \xrightarrow{k} Y$.

The next two lemmas will lead to the proof of Lemma 2 1.

Lemma 10 1 *Let $A\sigma B \xrightarrow{1} AB$ be a 1-sequence for positive A, B .*

We note the line $A\sigma B$ by L_1 and the other one AB by L_0 .

Then there exist positive A', B' , a generator σ' such as

$$A\sigma B = A'\sigma'B'$$

$$AB = A'B'$$

$$L_1 = A\sigma B = A'\sigma'B' \xrightarrow{1} L_0 = AB = A'B'$$

and $\text{rgcd}(L_1, L_0) = B'$.

(In the same way, we have $A\sigma B = A''\sigma''B''$ and $AB = A''B''$ for positive A'', B'' such as $A'' = \text{lgcd}(L_1, L_0)$).

PROOF:

We note $\text{rgcd}(L_1, L_0) = r$ and obviously $r \succ B$: $r = uB$ for a positive u .

We also have $A\sigma = xu$, $A = yu$ for positive x, y and

$$yu\sigma = xu$$

By Lemma 3 1, $y = y'd$ and $x = x'\sigma'd$ with $d = d_u^r(\sigma)$:
 $L_1 = A\sigma B = x'\sigma'duB$ and $L_0 = AB = y'duB$.
 Theses last give $(L_1, L_0) \succ duB$ and $uB = r \succ duB$.
 Finally $d = 1$ and we put $A' = y', B' = uB$ to obtain the result. (It's the same for lgcd).

In a more general case, we obtain the next result.

Lemma 11 1 *Let $L_k = A_1\sigma_{i_1}A_2\sigma_{i_2}A_3\dots A_k\sigma_{i_k}A_{k+1} \xrightarrow{1} L_{k-1} = A_1\sigma_{i_1}A_2\sigma_{i_2}A_3\dots A_kA_{k+1} \xrightarrow{k-2} L_1 = A_1\sigma_{i_1}A_2\dots A_kA_{k+1} \xrightarrow{1} L_0 = A_1A_2\dots A_kA_{k+1}$ be a k -sequence.*

Each line L_j , $0 \leq j \leq k$, satisfies $L_j = A_1\sigma_{i_1}A_2\sigma_{i_2}\dots\sigma_{i_{j-1}}A_j\sigma_{i_j}A_{j+1}\dots A_kA_{k+1}$ where σ_{i_r} for $j+1 \leq r \leq k$ are cancelled.

This sequence can be written in an equivalent way,

$$L_k = A'_1\sigma'_{i_1}A'_2\sigma'_{i_2}A'_3\dots A'_k\sigma'_{i_k}A'_{k+1} \xrightarrow{1} L_{k-1} = A'_1\sigma'_{i_1}A'_2\sigma'_{i_2}A'_3\dots A'_kA'_{k+1} \xrightarrow{k-2} L_1 = A'_1\sigma'_{i_1}A'_2\dots A'_kA'_{k+1} \xrightarrow{1} L_0 = A'_1A'_2\dots A'_kA'_{k+1}$$

such as $\text{rgcd}(L_k, \dots, L_1, L_0) = A'_{k+1}$.

PROOF:

Let $L_k = A_1\sigma_{i_1}A_2\sigma_{i_2}A_3\dots A_k\sigma_{i_k}A_{k+1} \xrightarrow{k} L_0 = A_1A_2\dots A_kA_{k+1}$ be a k -sequence.

We say this last satisfies (E) if $L_k = A'_1\sigma'_{i_1}A'_2\sigma'_{i_2}A'_3\dots A'_k\sigma'_{i_k}A'_{k+1} \xrightarrow{k} L_0 = A'_1A'_2\dots A'_kA'_{k+1}$ with $\text{rgcd}(L_k, \dots, L_0) = A'_{k+1}$.

We use here a double induction. Induction on k and on $l = \text{length}(L_0)$.

$E(k, l)$ is true, means that a k -sequence with $\text{length}(L_0) = l$, satisfies (E) as above.

The basis of induction are as follow:

- 1- $E(k', 0)$ is true for any k' . Indeed, any k' -sequence with $\text{length}(L_0) = 0$ gives $A_1 = \dots = A_{k+1} = 1$. Obviously, $\text{rgcd}(L_k, \dots, L_1, L_0 = 1) = 1 = A_{k+1}$. Here, the A'_i 's can be taken equals to $A_i = 1$'s.
- 2- $E(1, l)$ is true for any l , by Lemma 10 1.

We can then suppose $E(k', l-1)$ is true for any k' and $E(k', l)$ is true for any $0 \leq k' \leq k-1$, and let us prove that

$E(k, l)$ is true.

We have the k -sequence $L_k = A_1\sigma_{i_1}A_2\sigma_{i_2}A_3\dots A_k\sigma_{i_k}A_{k+1} \xrightarrow{k} L_0 = A_1A_2\dots A_kA_{k+1}$. We note $R = \text{rgcd}(L_k, \dots, L_0)$.

We consider the $(k-1)$ -sequence $L_{k-1} = A_1\sigma_{i_1}A_2\sigma_{i_2}A_3\dots\sigma_{i_{k-1}}A_kA_{k+1} \xrightarrow{k-1} L_0 = A_1A_2\dots A_kA_{k+1}$.

As $E(k-1, l)$ is true by hypothesis of induction, we have

$$L_{k-1} = A'_1\sigma'_{i_1}A'_2\sigma'_{i_2}A'_3\dots\sigma'_{i_{k-1}}A'_k \xrightarrow{k-1} L_0 = A'_1A'_2\dots A'_k$$

with $\text{rgcd}(L_{k-1}, \dots, L_0) = A'_k$.

Moreover, $L_j \succ A_k A_{k+1}$ for $0 \leq j \leq k-1$, so $\text{rgcd}(L_{k-1}, \dots, L_0) = A'_k \succ A_k A_{k+1}$. And $A'_k = V A_k A_{k+1}$ for a positive V .

We can finally write

$$L_k = A'_1 \sigma'_{i_1} A'_2 \sigma'_{i_2} A'_3 \dots \sigma'_{i_{k-1}} V A'_k \sigma_{i_k} A_{k+1}.$$

Now, if we write $A'_1 = \sigma B$, we obtain the following $k+1$ -sequence

$$L_k = \sigma B \sigma'_{i_1} A'_2 \sigma'_{i_2} A'_3 \dots \sigma'_{i_{k-1}} V A_k \sigma_{i_k} A_{k+1} \xrightarrow{k} L_0 = \sigma B A'_2 \dots A'_{k-1} V A_k A_{k+1} \xrightarrow{1} L'_0 = B A'_2 \dots A'_{k-1} V A_k A_{k+1}.$$

$R = \text{rgcd}(L_k, \dots, L_0)$ is such as $A'_k = V A_k A_{k+1} = \text{rgcd}(L_k, \dots, L_0) \succ R$, so $L'_0 \succ R$ and R common to $L_k, \dots, L_0, L'_0$ satisfies $\text{rgcd}(L_k, \dots, L_0, L'_0) = R$.

We deal with a $k+1$ -sequence such as $\text{length}(L'_0) = l-1$; as $E(k+1, l-1)$ is true, this last sequence can be written,

$$L_k = A''_0 \sigma'' A''_1 \sigma''_{i_1} A''_2 \sigma''_{i_2} A''_3 \dots \sigma''_{i_{k-1}} A''_k \sigma''_{i_k} R \xrightarrow{k} L_0 = A''_0 \sigma'' A''_1 A''_2 \dots A''_{k-1} A''_k R \xrightarrow{1} L'_0 = A''_0 A''_1 A''_2 \dots A''_{k-1} A''_k R.$$

Finally, by considering $L_k \xrightarrow{k} L_0$ with the above A''_i 's, our initial sequence is written in the right way.

We give now the proof of Lemma 2 1.

PROOF Lemma 2 1:

For $b = \Delta^{-r} b_+$ quasipositive, with algebraic length k , we have the k -sequence $b_+ \xrightarrow{k} \Delta^r$.

Lemma 7 1 gives $d_1, d_2, \dots, d_k$ such as

$$b_+ d_1 d_2 \dots d_k = \Delta^r d_1 \sigma_{i_1} d_2 \sigma_{i_2} \dots d_k \sigma_{i_k}.$$

Let s be an integer as large as needed such as $d_1 d_2 \dots d_k \prec \Delta^s$ and $r \leq s$.

We have, with $d_1 d_2 \dots d_k U = \Delta^s$, for a positive U , $b_+ \Delta^s = b_+ d_1 d_2 \dots d_k U = \Delta^r d_1 \sigma_{i_1} d_2 \sigma_{i_2} \dots d_k \sigma_{i_k} U$ by Lemma 7 1.

We can suppose s even, and $b_+ \Delta^s = \Delta^s b_+ = \Delta^r \Delta^{s-r} b_+$ and,

$$\Delta^{s-r} b_+ = d_1 \sigma_{i_1} d_2 \sigma_{i_2} \dots d_k \sigma_{i_k} U \xrightarrow{k} \Delta^{s-r} \Delta^r = d_1 d_2 \dots d_k U, (E)$$

In $b_+ \xrightarrow{k} \Delta^r$, each line of the sequence is noted by K_i : $b_+ = K_k$, $\Delta^r = K_0$. If we note $\Delta^{s-r} K_i = L_i$,

(E) as above gives $L_k = \Delta^{s-r} b_+ = d_1 \sigma_{i_1} d_2 \sigma_{i_2} \dots d_k \sigma_{i_k} U$, $L_0 = \Delta^{s-r} \Delta^r = d_1 d_2 \dots d_k U$.

Finally, each L_i is right divided by Δ^{s-r} and Lemma 11 1 gives, by supposing $s-r$ even,

$$\Delta^{s-r} b_+ = b_+ \Delta^{s-r} = A_1 \delta_{i_1} A_2 \delta_{i_2} \dots A_k \delta_{i_k} A_{k+1} \Delta^{s-r} \xrightarrow{k} \Delta^r \Delta^{s-r} = A_1 A_2 \dots A_k A_{k+1} \Delta^{s-r}$$

By simplifying by Δ^{s-r} , we have the result.

We give now the proof of Lemma 3 1.

PROOF Lemma 3 1:

Let $b = \Delta^r b_+$ be a quasipositive word and $b_+ = A_1 \sigma_{i_1} A_2 \sigma_{i_2} \dots A_k \sigma_{i_k} A_{k+1} \xrightarrow{k}$
 $\Delta^r = A_1 A_2 \dots A_k A_{k+1}$ be the corresponding k-sequence.

We have $b_+ = b_1 b_2 \dots b_r \dots b_s$ in its left normal form. If A_1 as above is written $A_1 = u_1 \dots u_p$ in its left normal form, we have:

$$1-A_1 \prec \Delta^r \text{ so } p \leq r \leq s$$

$$2-A_1 = u_1 \dots u_p \prec b_1 \dots b_p$$

We prove now that $A_1 = b_1 \dots b_p$.

By induction on k .

If $k = 1$. We deal with $b_+ = A_1 \sigma_{i_1} A_2$ and $\Delta^r = A_1 A_2$.

In one side, we have $A_1 \prec b_1 \dots b_p$.

In the other side, Lemma 10 1 tells us we can consider A_1 equals to $\text{lgcd}(b_+, \Delta^r)$. As, $p \leq r$, $b_1 \dots b_p \prec \Delta^r$: so, $b_1 \dots b_p \prec \text{lgcd}(b_+, \Delta^r) = A_1 \prec b_1 \dots b_p$.

Finally, $A_1 = b_1 \dots b_p$.

For any k . We suppose the result true for $k - 1$: for integers r, s, p ,

with $p \leq r \leq s$, such as $b_1 b_2 \dots b_r \dots b_s = C_1 \sigma_{i_1} C_2 \sigma_{i_2} \dots C_{k-1} \sigma_{i_{k-1}} C_k \xrightarrow{k-1}$
 $\Delta^r = C_1 C_2 \dots C_k$ and $C_1 = u_1 u_2 \dots u_p$ in its normal form, we have $C_1 = b_1 b_2 \dots b_p$.

Let $b_+ = b_1 b_2 \dots b_r \dots b_s = A_1 \sigma_{i_1} A_2 \sigma_{i_2} \dots A_k \sigma_{i_k} A_{k+1} \xrightarrow{k} \Delta^r = A_1 A_2 \dots A_k A_{k+1}$
be a k-sequence with $A_1 = u_1 u_2 \dots u_p \prec b_1 b_2 \dots b_p$.

Now, as $p \leq r$, $\Delta^r = b_1 b_2 \dots b_p U b_{t+1} \dots b_s$ for a positive U and $s + p - r = t$.

The sequence $b_+ = b_1 b_2 \dots b_p b_{p+1} \dots b_t b_{t+1} \dots b_s \xrightarrow{k} \Delta^r = b_1 b_2 \dots b_p U b_{t+1} \dots b_s$
is equivalent to

$$b_{p+1} \dots b_t \xrightarrow{k} U$$

we can write as $b_{p+1} \dots b_t = X_k \sigma_{i_k} Y_k \xrightarrow{1} X_k Y_k \xrightarrow{k-1} U$.

By lemmas above, we obtain

$$b_1 b_2 \dots b_p b_{p+1} \dots b_t b_{t+1} \dots b_s = b_1 b_2 \dots b_p X_k \sigma_{i_k} Y_k b_{t+1} \dots b_s = A_1 \sigma_{i_1} A_2 \sigma_{i_2} \dots A_k \sigma_{i_k} A_{k+1} \xrightarrow{1}$$

$$b_1 b_2 \dots b_p X_k Y_k b_{t+1} \dots b_s = A_1 \sigma_{i_1} A_2 \sigma_{i_2} \dots A_k A_{k+1} \xrightarrow{k-1} b_1 b_2 \dots b_p U b_{t+1} \dots b_s.$$

Finally the $(k-1)$ -sequence $L_{k-1} = b_1 b_2 \dots b_p X_k Y_k b_{t+1} \dots b_s = A_1 \sigma_{i_1} A_2 \sigma_{i_2} \dots A_k A_{k+1} \xrightarrow{k-1}$
 $L_0 = b_1 b_2 \dots b_p U b_{t+1} \dots b_s$ with $A_1 \prec b_1 b_2 \dots b_p$ gives by induction

$$A_1 = b_1 b_2 \dots b_p.$$

Moreover, in the previous sequence, we can notice that by Lemma 11 1

$\text{rgcd}(L_k, L_{k-1}, \dots, L_0) \succ b_{t+1} \dots b_s$; as we can suppose $A_{k+1} = \text{rgcd}(L_k, L_{k-1}, \dots, L_0)$,
 $A_{k+1} = A'_{k+1} b_{t+1} \dots b_s$ for a positive A'_{k+1} .

We can now conclude by

$$b_{p+1} \dots b_t = \sigma_{i_1} A_2 \sigma_{i_2} \dots A_k \sigma_{i_k} A'_{k+1} \xrightarrow{k} U = A_2 A_3 \dots A_k A'_{k+1}.$$

In the next paragraph, we give a computational algorithm and its complexity.

Algorithm We give now the practical consequence of Lemma 11 1.

Lemma 12 1 *Let $b = \Delta^{-r}b_1b_2...b_r...b_s$ be a quasipositive word. By previous results, for p such as $1 \leq p \leq r \leq s$, there exist k positive words $A_2, A_3, ..., A_k, A_{k+1}$ and k generators such as*

$$b_{p+1}...b_{p+s-r} = \sigma_{i_1}A_2\sigma_{i_2}A_3...A_k\sigma_{i_k}A_{k+1}$$

$$U = A_2A_3...A_kA_{k+1}$$

where U satisfies $b_1b_2...b_pUb_{p+s-r+1}...b_s = \Delta^r$.
Then, we can write

$$b_{p+1}...b_{p+s-r} = \sigma'_{i_1}A'_2\sigma'_{i_2}A'_3...A'_k\sigma'_{i_k}A'_{k+1}$$

$$U = A'_2A'_3...A'_kA'_{k+1}$$

such as , for $2 \leq j \leq k+1$, $A'_j = \text{lgcd}(A'_j\sigma'_{i_j}A'_{j+1}...A'_k\sigma'_{i_k}A'_{k+1}, A'_jA'_{j+1}...A'_kA'_{k+1})$

PROOF:

If for a k -sequence $\sigma_{i_1}A_2\sigma_{i_2}A_3...A_k\sigma_{i_k}A_{k+1} \xrightarrow{k} A_2A_3...A_kA_{k+1}$, there exist A'_j as in the lemma above, we say that the sequence satisfies (E).

We use here a double induction. Induction on k and on $l = \text{length}(A_2...A_{k+1})$. We say that $E(k, l)$ is true if any k -sequence of length l satisfies (E).

The basis of induction are:

1- $E(k, 0)$ true for any k . Indeed, $l = 0$ means $A_j = 1$ and the k -sequence is $\sigma_{i_1}\sigma_{i_2}...\sigma_{i_k} \xrightarrow{k} 1$. Obviously, $A_j = 1 = \text{lgcd}(\sigma_{i_1}\sigma_{i_2}...\sigma_{i_k}, 1)$.

2- $E(1, l)$ true for any l . The 1-sequence is $\sigma_{i_1}A_2 \xrightarrow{1} A_2$ and obviously, $A_2 = \text{lgcd}(A_2, A_2)$.

We can suppose $E(k, l-1)$ true for any k and $E(k', l)$ true for $1 \leq k' \leq k-1$ and let us prove $E(k, l)$.

Let A_d be the first factor not equal to 1, the k -sequence becomes $\sigma_{i_1}...\sigma_{i_d}A_d...A_k\sigma_{i_k}A_{k+1} \xrightarrow{k} A_d...A_{k+1}$.

As $A_d = \delta A'_d$, we get a $(k+1)$ -sequence $X = \sigma_{i_1}...\sigma_{i_d}\delta A'_d...A_k\sigma_{i_k}A_{k+1} \xrightarrow{k} Y = A'_d...A_{k+1}$.

We have $\text{length}(A'_d...A_k) = l-1$ and as $E(k, l-1)$ true for any k ,

$$X = C_1\delta_{i_1}C_2\delta_{i_2}C_3...C_{k+1}\delta_{i_{k+1}}C_{k+2}$$

$$Y = C_1C_2...C_{k+1}C_{k+2}$$

Deleting δ corresponds to deleting δ_{i_d} so $\delta A'_d \dots A_{k+1} = \delta Y = \delta C_1 C_2 \dots C_{k+1} C_{k+2} = C_1 \dots C_d \delta_{i_d} C_{d+1} \dots C_{k+2}$.

1-If $C_1 = \alpha C'_1$ is not equal to one, from the k -sequence of length l , $\sigma_{i_1} C_1 \delta_{i_1} C_2 \delta_{i_2} C_3 \dots C_{k+1} \delta_{i_{k+1}} C_{k+2} \xrightarrow{k} C_1 C_2 \dots C_{k+1} C_{k+2}$, we extract the other one of length $(l-1)$ where all generators, but δ_{i_d} , are deleted

$$X' = \sigma_{i_1} C'_1 \delta_{i_1} C_2 \delta_{i_2} C_3 \dots C_{k+1} \delta_{i_{k+1}} C_{k+2} \xrightarrow{k} Y' = C'_1 C_2 \dots C_{k+1} C_{k+2}.$$

$E(k, l-1)$ is true so $X' = D_1 \delta'_{i_1} D_2 \delta'_{i_2} D_3 \dots D_k \delta'_{i_k} D_{k+1}$ and $Y' = D_1 D_2 \dots D_{k+1}$ with

$$D_j = \text{lgcd} (D_j \delta'_{i_j} D_{j+1} \dots D_k \delta'_{i_k} D_{k+1}, D_j D_{j+1} \dots D_k D_{k+1}).$$

Moreover, $\text{lgcd} (X, Y) = \text{lgcd} (\alpha X', \alpha Y') = \alpha D_1$ and we obtain all datas for the final result, (F).

2-If $C_1 = 1$. Then $\text{lgcd} (X, Y) = 1$. We note $R = \text{lgcd} (X, \delta Y)$ and let α such as $\alpha \prec R$.

2.1-if α different from δ , then $\alpha \prec (X, \delta Y)$ gives $\alpha \prec (X, Y)$. As $\text{lgcd} (X, Y) = 1$, $\alpha = 1$.

$$2.2\text{-if } \alpha = \delta, \delta \prec X = \delta_{i_1} C_2 \delta_{i_2} C_3 \dots C_{k+1} \delta_{i_{k+1}} C_{k+2}.$$

$$2.2.1\text{-if } \delta = \delta_{i_1} \text{ then } X = \delta C_2 \delta_{i_2} C_3 \dots C_{k+1} \delta_{i_{k+1}} C_{k+2} \text{ and } Y \delta Y = \delta C_2 \dots C_{k+1} C_{k+2}.$$

As before, for $\sigma_{i_1} C_2 \delta_{i_2} C_3 \dots C_{k+1} \delta_{i_{k+1}} C_{k+2} \xrightarrow{k} C_2 \dots C_{k+1} C_{k+2}$, $E(k, l-1)$ being true, we have the final result, see (F).

2.2.2-if δ is different from δ_{i_1} then $\delta \prec X = \delta_{i_1} C_2 \delta_{i_2} C_3 \dots C_{k+1} \delta_{i_{k+1}} C_{k+2}$ gives $\delta \prec X' = C_2 \delta_{i_2} C_3 \dots C_{k+1} \delta_{i_{k+1}} C_{k+2}$. Moreover, $\delta \prec \delta Y = C_2 \dots C_d \delta_{i_d} C_{d+1} \dots C_{k+2}$.

As $E(k-1, l)$ is true, the $(k-1)$ -sequence being $X' = C_2 \delta_{i_2} C_3 \dots C_{k+1} \delta_{i_{k+1}} C_{k+2} \xrightarrow{k-1} \delta Y$

where all δ_{i_j} but δ_{i_d} are deleted, we obtain

$$X' = \delta D_1 \delta'_{i_1} D_2 \delta'_{i_2} D_3 \dots D_{k-1} \delta'_{i_k} D_k$$

$$\delta Y = \delta D_1 \dots D_k$$

We use now Lemma 11.1. Given a k -sequence $L_k = A_1 \sigma_{i_1} A_2 \sigma_{i_2} A_3 \dots A_k \sigma_{i_k} A_{k+1} \xrightarrow{k} L_0 = A_1 \dots A_{k+1}$

where σ_{i_k} is first deleted, then $\sigma_{i_{k-1}}$... and so on, we can suppose $\text{rgcd} (L_k, \dots, L_0) = A_k$.

By symmetry, if we delete first σ_{i_1} , then σ_{i_2} ... and so on, we can suppose $A_1 = \text{lgcd} (L_k, \dots, L_0)$.

Now, if we consider $L_k = \delta_{i_1} X' = \delta_{i_1} \delta D_1 \delta'_{i_1} D_2 \delta'_{i_2} D_3 \dots D_{k-1} \delta'_{i_k} D_k \xrightarrow{1} L_{k-1} = \delta D_1 \delta'_{i_1} D_2 \delta'_{i_2} D_3 \dots D_{k-1} \delta'_{i_k} D_k \xrightarrow{1} L_{k-2} = \delta D_1 D_2 \delta'_{i_2} D_3 \dots D_{k-1} \delta'_{i_k} D_k \xrightarrow{1} L_{k-3} = \delta D_1 D_2 D_3 \delta'_{i_3} \dots D_{k-1} \delta'_{i_k} D_k \xrightarrow{k-3} L_0 = \delta D_1 D_2 D_3 \dots D_{k-1} D_k$, we can write L_k as

$$L_k = \delta E_1 \sigma''_{i_1} E_2 \sigma''_{i_2} E_3 \dots E_k \sigma''_{i_k} E_{k+1}$$

$$L_0 = \delta E_1 E_2 E_3 \dots E_k E_{k+1}$$

Finally, $X = L_k$ and $Y = L_0$ and doing as in (F), with $\sigma_{i_1} E_1 \sigma''_{i_1} E_2 \sigma''_{i_2} E_3 \dots E_k \sigma''_{i_k} E_{k+1} \xrightarrow{k} E_1 E_2 E_3 \dots E_k E_{k+1}$, we obtain the final result.

REMARK:

This last result gives practical consequences for the next algorithm. Indeed, with the following datas,

$$P \xrightarrow{k} N$$

$$P = \sigma_{i_1} A_2 \sigma_{i_2} A_3 \dots A_k \sigma_{i_k} A_{k+1}$$

$$N = A_2 A_3 \dots A_k A_{k+1}$$

σ_{i_1} is easy to compute, among the generators which begin P .
Moreover A_2 satisfies $A_2 = \text{lgcd}(\sigma_{i_1}^{-1} P, N) \dots$ and so on.

Theorem 2 1 ALGORITHM

Let b be a word in B_{n+1} . For a positive word w , we note $\text{beg}(w)$ the set of generators left dividing w , $\text{beg}(w) = (\sigma / \sigma \prec w)$.

- 1- We compute the algebraic length of b , k :
if $k \leq 0$, STOP (b is not quasipositive; if $k = 0$, b is quasipositive iff $b = 1$)
elsewhere we go to step 2
- 2- We compute the left normal form of b , $b = \Delta^{-r} b_1 \dots b_r b_{r+1} \dots b_s$:
if $r \leq 0$, STOP (b is positive so quasipositive)
elsewhere we go to step 3
- 3- For each $0 \leq j \leq r$, we compute $P_j = b_{j+1} \dots b_{s-r+j}$ and we compute N_j such as $b_1 b_2 \dots b_j N_j b_{s-r+j+1} \dots b_s = \Delta^r$
- 4- We consider the set S with $S = [P_0, \dots, P_r, N_0, \dots, N_r]$ and the empty set S'
- 5- We begin with $j = 0$
- 6- We set $P = P_j$ and $N = N_j$ and we add P to the set S'

7-We set $m = 1$

8-For all elements P in the set S' :

we compute $\text{beg}(P)$ and for each σ in $\text{beg}(P)$, we compute P' such as $P = \sigma P'$.

For each P' , we compute the normal form of $N^{-1}P'$, we can write as $N^{-1}P' = A^{-1}B$.

We add all theses P', A, B in the set S and we add all the B in the set S' .

10-we set $m := m + 1$

9-If $m = k$, we set $j := j + 1$:

if $j = r + 1$ then we go to step 10

elsewhere, we go to step 6

Elsewhere, if $m \leq k - 1$, we go to step 8.

10-If 1 belongs to the set S , then b is quasipositive. Keeping trace to the factors used, by noting them by $A_2, \dots, A_{k+1}$ and $\sigma_{i_1}, \dots, \sigma_{i_k}$, we output the quasipositive form of a conjugate of b :

$$b' = A_{k+1}^{-1} \dots A_2^{-1} \sigma_{i_1} A_2 \sigma_{i_3} \dots \sigma_{i_k} A_{k+1}$$

and the form of b , $b = T_j^{-1} b' T_j$ with $T_j = (b_{s-r+j+1} \dots b_s)$.

Elsewhere, STOP (b is not quasipositive).

EXAMPLES:

In B_4 , we put $b = 2212^{-1}2^{-1}2^{-1}2^{-1}322$ with $k = 2$; its normal form is

$$b = \Delta^{-4} 2321.32123.12321.32123.12.2132.2$$

We have $r = 4$, $s = 6$ and $b_1 = 2321$, $b_2 = 32123$, $b_3 = 13221$, $b_4 = 12$, $b_5 = 2132$, $b_6 = 2$

so $a_1 = 23$, $a_2 = 2$, $a_3 = 2$, $a_4 = 1321$, $a_5 = 13$, $a_6 = 13213$.

We have $P = b_3 b_4 b_5 = 12321.32123.12$; the obtained N is $N = a_2 \bar{a}_1 a_6 \bar{a}_5 = 2.21.13213.31 = 2.21.132113.3$.

But $P = 123231212312 = 122321212312 = 122312112312 = 122132112132 = 122132121232 = 1.2213.1211232 = 1.221132113.2.3$.

We have $P = 1.A_2.2.A_3$ with $A_2 = 221132113$ and $A_3 = 3$: as $N = A_2 A_3$ b is quasipositive.

If we consider now $b = 123^{-1}$ with $k = 1$; its normal form is

$b = \Delta^{-1} 323.2132$ so $b_1 = 323$, $b_2 = 2132$ and $r = 1$; and $a_1 = 123$, $a_2 = 13$.

We have two possibilities for P :

$P = b_2 = 2132$ and $N = a_1 = 123$: P can't be reduced to N

$P = b_1 = 323$ and $N = \bar{a}_2 = 31$: P can't be reduced to N

Finally b is not quasipositive.

If we consider now b , a word of algebraic length k written in its left normal form $b = \Delta^{-r}b_+$, and we note by l the length of b_+ , we obtain the next result.

Theorem 3 1 *The complexity of the previous algorithm is $O(n^k l^2)$.*

PROOF:

For the initial P_j and N_j , we have $r + 1$ choices. Let us note by x the complexity for computing the normal form of a word.

At the end of the algorithm, the set S contains at most $(r+1)n^k$ elements: for an initial P_j , we have to compute $beg(P)$ k times and we have n choices of generators for this last.

Moreover at each step, we have to compute the normal form of the elements we are dealing with (step 8), so the final complexity is about

$$(r + 1)n^k x.$$

In different litterature [3], it's proved that x is about nl^2 .

Finally, the final complexity is at most $n^{k+1}l^2$, which is $O(n^k l^2)$.

Conclusion This algorithm and all its proofs only need the combinatorial properties of braid groups: the existence of $lgcd$, $rgcd$ and Δ , and as a consequence, the existence of a normal form for every word and over all, the existence of a length function. Theses conditions beeing the classical definition of Garside groups, as braid groups, spherical Artin-Tits groups, (see [2],[3],[4]), the previous algorithm can then be used in all theses groups, with a same complexity.

References [1]BIRMAN, Joan S. Braids, links, and mapping class groups. *Annals of Mathematics Studies*, No. 82. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1974.

[2]BRIESKORN,SAITO. Artin gruppen und Coxeter gruppen. *Invent. Math.* 17, 1972, 245-271

[3]DEHORNOY. Groupes de Garside. *Ann. Sc. Ec. Norm. Sup.*, No. 35, 2002, 267-306

[4]DEHORNOY,PARIS. Gaussian groups and Garside groups, two generalisations of Artin groups. *Proc. London. Math. Soc.*(3), No 79, 1999, 569-604.

[5]GARSIDE, F. A. The braid group and other groups. *Quart. J. Math. Oxford Ser+.* (2) 20 1969 235-254.

[6]EL-RIFAI, ELSAYED A.; MORTON, H. R. Algorithms for positive braids. *Quart. J. Math. Oxford Ser.* No 45 (1994), no. 180, 479-497.

[7]OREVOV, S. Yu. Erratum to: "Link theory and oval arrangements of real algebraic curves" [*Topology* 38 (1999), no. 4, 779-810; MR 2000b:14066].

Topology 41 (2002), no. 1, 211–212. 14P25 (20F36 57M25). 34 (2000), no. 3, 84–87.

[8]OREVKOV, S. Yu. Quasipositivity test via unitary representations of braid groups and its applications to real algebraic curves. *J. Knot Theory Ramifications* 10 (2001), no. 7, 1005–1023.

[9]RUDOLPH, Lee Quasipositive pretzels. *Topology Appl.* 115 (2001), no. 1, 115–123.

[10]RUDOLPH, Lee Quasipositive plumbing (constructions of quasipositive knots and links. V). *Proc. Amer. Math. Soc.* 126 (1998), no. 1, 257–267.

A.Bentalha

Universite Paul abatier

Laboratoire Emile Picard, 31400 Toulouse, France

a.bentalha@laposte.net